

College algebra  
Direct, Inverse, and Joint Variation  
Section 2.5

If two variables are related to each other by a constant ratio or product, we say we have a direct or inverse variation.

**Direct Variation:**

We will start with an example.

Margie gets paid \$10 per hour when she babysits. Fill in the table for the various number of hours given.

x, number of hours	y, total charge, dollars	Find $y/x$
1	\$10	$10/1 = 10$
2	\$20	$20/2 = 10$
3	\$30	$30/3 = 10$
7	\$70	$70/7 = 10$

The ratio is constant which means there is a direct variation here.

The total charge varies directly as the number of hours worked.

$$y = 10x$$

**Definition: Direct Variation:**

If a situation can be modeled by the linear function  $f(x) = kx$ , or  $y = kx$ , where  $k$  is a nonzero constant, we say that it is a direct variation. We could say  $y$  varies directly as  $x$  or  $y$  is directly proportional to  $x$ . The number  $k$  is the variation constant or the constant of proportionality.

Can you think of any other variables that would be directly proportional?

$x = \#$  of lemon cakes bought

$y = \text{total cost}$

if cakes are \$7 each

$$y = 7x$$

## Inverse Variation:

Again, let's look at an example. Margie visits her aunt who lives 50 miles away. Fill in the table for the various modes of transport given.

$r$ , rate of travel, mph	$t$ , time, hours	Find $r \cdot t$
50 (She drives.)	1	$50 \cdot 1 = 50$
10 (She bikes.)	5	$10 \cdot 5 = 50$
5 (She walks.)	10	$5 \cdot 10 = 50$
1 (She crawls.)	50	$1 \cdot 50 = 50$

The product is constant which means there is an inverse variation here.

$$d = r \cdot t$$

$d$  = distance  
 $r$  = rate  
 $t$  = time

Margie's rate is inversely proportional to her time.

### Definition: Inverse Variation:

If a situation can be modeled by the linear function  $f(x) = k/x$ , or  $y = k/x$ , where  $k$  is a nonzero constant, we say that it is an inverse variation. We could say  $y$  varies inversely as  $x$  or  $y$  is inversely proportional to  $x$ . The number  $k$  is the variation constant or the constant of proportionality.

Can you think of any other variables that would be inversely proportional?

For Margie, the  $k$  is the constant distance to Auntie's.

expl 1: Find the variation constant and the equation of variation for the given situation.

a.)  $y$  varies directly as  $x$ , and  $y = 54$  when  $x = 12$

$$y = kx$$

$$\frac{54}{12} = \frac{k \cdot 12}{12}$$

$$4.5 = k$$

var. const

$$y = kx$$

$$y = 4.5x$$

eqn of var.

b.)  $y$  varies *inversely* as  $x$ , and  $y = 12$  when  $x = 5$

$$y = k/x$$

$$12 = k/5$$

$$60 = k$$

var const

$$y = k/x$$

$$y = 60/x$$

eqn of var.

$W$  = weight that a hor. beam can support (kg)  
 $L$  = length of beam (m)

expl 2: The weight  $W$  that a horizontal beam can support varies inversely as the length  $L$  of the beam. Suppose an 8 meter beam can support 1200 kg. How many kilograms can a 14 meter beam support?

$$W = k/L$$

$$1200 = k/8$$

$$9600 = k$$

$$\therefore W = 9600/L$$

$$W = 9600/14$$

$$W \approx 686 \text{ kg.}$$

A 14-m beam can support 686 kg approx.

expl 3: The relative aperture, or f-stop, of a 23.5-mm diameter camera lens is directly proportional to the focal length  $F$  of the lens. If a focal length of 150 mm has an f-stop of 6.3, find the f-stop of this lens with a focal length of 80 mm.

$f$  = f-stop, relative aperture

$F$  = focal length of lens (mm)

$$f = k \cdot F$$

$$\frac{6.3}{150} = \frac{k \cdot 150}{150}$$

$$k = 0.042$$

$$f = 0.042 \cdot F$$

$$= 0.042 (80)$$

$$f = 3.36 \text{ f-stop}$$

$$y = kx \text{ OR } y = k/x?$$

Find  $k$  and then form the equation of variation.

$$y = kx \text{ OR } y = k/x?$$

Find  $k$  and then form the equation of variation.



### Combined Variation:

There are three related types of variation we will study.

1.  $y$  varies directly as the  $n$ th power of  $x$  if there is some nonzero constant  $k$  such that

$$y = kx^n.$$

2.  $y$  varies inversely as the  $n$ th power of  $x$  if there is some nonzero constant  $k$  such that

$$y = \frac{k}{x^n}.$$

3.  $y$  varies jointly as  $x$  and  $z$  if there is some nonzero constant  $k$  such that  $y = kxz$ .

### General Procedure:

1. Read what type of variation (direct, inverse, joint, or some combination) the variables have in the problem.
2. Start with the general equation given above. Define your variables!
3. Use the given information to find the missing proportionality constant  $k$ .
4. Rewrite the equation with your value of  $k$  in place. Use it to answer the question.

expl 4: The weight  $W$  of an object varies inversely as the square of the distance  $d$  from the center of the earth. At sea level (3978 miles from the center of the earth), an astronaut weighs 220 pounds. Find his weight when he is 200 miles above the surface of the earth.

$W$  = weight of object

$d$  = distance from center of Earth

$$W = k/d^2$$

$$220 = k / 3978^2$$

$$k = 3,481,386,480$$


$$W = \frac{3,481,386,480}{d^2}$$

$$W = \frac{3,481,386,480}{4178^2}$$

$$W \approx 199.44 \text{ lbs}$$

$$W = \frac{k}{d^2}$$

What is  $d$  when he is 200 miles above the earth?



4178 miles.

Pg 2  $y = k/x$

expl 5: Find the variation constant and the equation of variation for the given situation.

y varies jointly as x and z and inversely as w, and y = 14 when x = 3, z = 2, and w = 2

$$y = \frac{kxz}{w}$$

$$y = \frac{14xz}{3w}$$

eqn of var.

$$14 = \frac{k \cdot 3 \cdot 2}{2}$$

$\frac{14}{3} = k$  var. constant

expl 6: The kinetic energy K of a moving object varies jointly with its mass m and the square of its velocity v. An object with a mass of 25 kilograms that is moving with a velocity of 10 meters per second has a kinetic energy of 1250 joules. If this same object is now moving at 35 meters per second, what is its kinetic energy?

K = Kinetic energy of object (joules)

m = mass of object (kg)

v = Velocity of object (m/s)

First, we find k, the proportionality constant. Do not confuse this with K, the kinetic energy.

$$K = k m v^2$$

$$1250 = k \cdot 25 \cdot 10^2$$

$$1250 = k \cdot 2500$$

$$\frac{1250}{2500} = k$$

$\frac{1}{2}$  or 0.5 = k

$$K = 0.5 m v^2$$

$$= 0.5 (25) \cdot 35^2$$

$$K = 15312.5 \text{ joules}$$



.....

.....

.....

.....

.....

.....



.....

.....

.....

.....

.....

.....

.....

