

We will use our log rules and the equivalency of $x = a^y$ and $y = \log_a x$.

College algebra

Class notes

Solving Exponential and Logarithmic Equations (section 6.6)

Recall: Definition: Exponential Equation: An equation with the variable in the exponent position will be called an exponential equation.

Examples:

$$5^x = 5^4, \quad e^{2t} = 500, \quad 3^{4x} = 81, \quad 3^r = 2^{r-1}, \quad e^x - 6e^{-x} = 1, \quad 27 = 3^{5x} \cdot 9^{x^2}$$

Recall: Definition: Logarithmic Equation: An equation that has logs of variable expressions will be called a logarithmic (or log) equation.


Examples:

$$\log_3(4w) = 4, \quad 3\log_2(x-1) + \log_2 4 = 5, \quad \log_4 x + \log_4(x-3) = 1, \quad \log_2(x-1) - \log_6(x+2) = 2$$

We're solving equations in these sections. Remember we are trying to find the x that makes the equation true. We will need to keep a lot in our minds.


- 1.) the logarithm rules from the previous section
- 2.) the fact that $x = a^y$ and $y = \log_a x$ are equivalent
- 3.) the two properties below

Base-Exponent Property: For any $a > 0$ and $a \neq 1$, we know that $a^x = a^y$ if and only if $x = y$.



So if $5^x = 5^4$, then what must be true?

Logarithmic Equality Property: For any $M > 0$, $N > 0$, $a > 0$, and $a \neq 1$, we know that $M = N$ if and only if $\log_a M = \log_a N$.



So if $\log_2 x = \log_2 10$, then what must be true?


Many of the problems can be solved by various methods. We will explore this with the examples. When doing homework, choose the method that most appeals to you or that best fits that equation.

Solving Exponential Equations:

expl 1: Solve. Try the different methods below.


$$4^{2t+1} = 20$$

Method 1: Use the equivalency of $x = a^y$ and $y = \log_a x$ to rewrite the equation in log form.




Method 2: Use the **Logarithmic Equality Property** to “take the log (base 4) of both sides”.

Method 3: Use the **Logarithmic Equality Property** to “take the natural log of both sides”.



expl 2: Solve.


$$3^{4x} = 81$$




If you know your
powers of 3, it might
get you going here.

expl 3: Solve.

$$2^{2x} + 2^x - 12 = 0$$



This looks crazy
but substitute y
for 2^x and ...




Always
check your
answers.

Solving Logarithmic Equations: Some equations will need to be simplified using our newly learned log rules in addition to using the exponential and logarithmic properties on page 1.

expl 4: Solve. Try the different methods below.


$$\log_3(4w) = 4$$

Method 1: Use the equivalency of $x = a^y$ and $y = \log_a x$ to rewrite the equation in exponential form.




Always
check your
answers.

Method 2: Use the **Base-Exponent Property** to rewrite this as an exponential equation. Then use the log rules to simplify as you solve for w .



What base
will you
use?




Can we get this to the
 $\log_a M = \log_a N$ form?

expl 5: Solve. Express irrational solutions in exact form.


$$2\log_2(x-1) = \log_2 5$$

expl 6: Solve.

$$\log_3(2x-5) = 1 - \log_3 x$$



Get the logs together and use your
log rules. Can you rewrite this as a
quadratic equation and solve?




The methods we use
sometimes produce
extraneous solutions. You
must check your solutions
in the *original* equation.

Solving Equations Graphically: As we have seen before, solving an equation graphically is simply a matter of graphing “ $y =$ the left side” and “ $y =$ the right side” and seeing where they intersect. One advantage of a graphical solution is that you *never* get extraneous solutions.

expl 7: Solve using a graphing calculator. Copy the graph here. Do *not* just TRACE. Use the INTERSECT function on the calculator. Round your solutions to three decimal places.


$$\log_3(2x - 5) = 1 - \log_3 x$$



You will need the Change of Base Formula. Do *not* use the other log rules.

expl 8: Solve using a graphing calculator. Copy the graph here. Do *not* just TRACE. Use the INTERSECT function on the calculator. Round your solutions to three decimal places.

$$2^x - 5 = 3x + 3$$



Did you find both solutions?

Optional Worksheet: Using log rules to solve equations:

This worksheet guides you through solutions with step-by-step instructions, providing practice solving equations both algebraically and graphically. It gives good advice on how to graph the pieces of these equations.

expl 9: The value of a Chevy Cruze LT (in dollars) that is t years old can be modeled by $V(t) = 19,200(0.82)^t$. Answer the following questions.

a.) How much is the car worth when $t = 0$? Interpret this result.

b.) When will the car be worth \$10,000? Round to the nearest tenth of a year.

